

Neural Network Related to Hand Movement: A Combined Study of Diffusion Tensor Tractography and Functional MRI

Ji HEON HONG, PT, MS¹⁾, SUNG HO JANG, MD¹⁾

¹⁾Department of Physical Medicine and Rehabilitation, College of Medicine, Yeungnam University:
317-1, Daemyungdong, Namku, Taegu, 705-717, Republic of Korea.

TEL: 82-53-620-3269, FAX: 82-53-620-3269, E-mail: strokerehab@hanmail.net, belado@med.yu.ac.kr

Abstract. [Purpose] Little is known about the detailed anatomical connections of the neural network related to hand movement in the human brain. We investigated the neural network using diffusion tensor tractography (DTT) data analyzed in conjunction with functional MRI (fMRI) activation results. [Subjects and Method] We recruited 19 healthy volunteers for this study. Probabilistic tractography was used to analyze diffusion tensor imaging (DTI) data that were collected using fMRI activation induced by grasp-release movements of the hand at a rate of 1 Hz. [Results] The brain areas connected to the primary sensorimotor cortex (SM1), which is activated by hand movements, were the premotor cortex (100%), superior parietal lobule (100%), intraparietal sulcus (100%), supramarginal gyrus (97.37%), supplementary motor area (89.47%), thalamus (86.84%), putamen (81.58%), pars opercularis (81.58%), pars triangularis (68.42%), angular gyrus (65.79%), and cerebellum (60.53%) in the same hemisphere and the contralateral primary motor cortex (60.53%) in the opposite hemisphere. No significant difference was observed in the total incidence of connected tracts between hemispheres. [Conclusion] These results reveal that more brain areas are involved in hand movements than were previously thought necessary for motor planning and execution in the human brain.

Key words: Motor control, Neural network, Diffusion tensor tractography

(This article was submitted Jun. 30, 2010, and was accepted Aug. 12, 2010)

INTRODUCTION

It is necessary to investigate the neural network to understand the physiological mechanisms of normal movement in the human brain¹⁾. Several brain evaluation techniques (e.g., functional neuroimaging, electroencephalography, magnetoencephalography) have been used to elucidate functional relationships between neural activity and movement control²⁻⁷⁾. Such studies have reported that various cortical and subcortical areas participate in movement control²⁻⁷⁾. However, these evaluation techniques are insufficient for resolving and visualizing the cortico-subcortical neural network for movement.

Diffusion tensor tractography (DTT), which is derived from diffusion tensor imaging (DTI), allows us to visualize and localize the architecture and integrity of subcortical neural tracts in three dimensions⁸⁻¹⁰⁾. Functional MRI (fMRI) is capable of precisely identifying cortical activation sites because of its high spatial resolution¹¹⁾. Therefore, when DTT data are analyzed in conjunction with fMRI results (activation in response to movement) we can obtain a more accurate characterization of the cortico-subcortical neural network related to movement¹²⁻¹⁴⁾.

In the current study we elucidated the neural network related to hand movements using DTT data analyzed in conjunction with fMRI activation results.

SUBJECTS AND METHOD

We recruited 19 right-handed normal subjects (17 male, 2 female; mean age: 28.9 years; range: 25–33 years) with no previous history of neurological, psychiatric, or physical illness. Handedness was evaluated using the Edinburgh Handedness Inventory¹⁵⁾. All subjects understood the purpose of this study, and they provided their written, informed consent prior to participation in this study. This study was approved by the institutional review board of our university hospital.

The blood oxygenation level dependent (BOLD) fMRI measurement, which employs the echo planar imaging (EPI) technique, was performed using a 1.5-T Philips Gyroscan Intera scanner (Hoffman-LaRoche, Ltd.; Best, the Netherlands) with a standard head coil. The EPI BOLD images were acquired over 20 identical axial sections, producing a total of 1,200 images for each subject. Imaging conditions were TR/TE = 2 sec/60 msec, FOV = 210 mm,




Fig. 1. (A) The contralateral primary sensorimotor cortex, which was activated by hand movements, was chosen as the seed region of interest (left). The probabilistic map of the neural network-related hand movements (right). (B) The quantitative network is color coded for 25 (dark red) to 5,000 (white) samples passing through the voxel. M1: primary motor cortex; SPL: superior parietal lobule; IPS: inferior parietal sulcus; SMA: supplementary motor area.

matrix size = 64×64 , and slice thickness = 5 mm. Subjects were examined supine with their eyes closed, and they were secured firmly with the forearm in supination. A repetitive alternating cycle of control (rest for 20 seconds) and stimulation (activity for 20 seconds) with grasp-release hand movements at a metronome-guided frequency of 1 Hz was performed for the motor task. Each “control and stimulation” task (40 seconds) was repeated three times.

The DTIs were acquired using a synergy-L Sensitivity-encoding (SENSE) head coil on a 1.5-T Philips Gyroscan Intera scanner (Hoffman-LaRoche, Ltd.; Best, the Netherlands) with a navigator echo. The DTI data were collected using a single-shot spin-echo planar imaging sequence of $1.73 \times 1.73 \times 2.3 \text{ mm}^3$. Sixty-seven contiguous slices (matrix = 128×128 , field of view = $221 \times 221 \text{ mm}^2$, repetition time/echo time = 10,726/76 ms, SENSE factor = 2, EPI factor = 67, $b = 1000 \text{ s/mm}^2$, NEX = 1, and thickness = 2.3 mm) were acquired for each of the 32 non-collinear diffusion-sensitizing gradients.

The fMRI data were analyzed using SPM2 software (Wellcome Department of Cognitive Neurology, UK) run in MATLAB (Mathworks, USA). The functional images were realigned by slice timing and motion corrected. The data were coregistered and resliced using the diffusion-weighted EPI volume with the highest signal-to-noise ratio (no diffusion weighting, $b = 0$) for each subject as a template¹⁶. The final data were smoothed with an 8 mm isotropic Gaussian kernel. Statistical parametric maps were obtained, and voxels were considered significant at a p-value of 0.05, corrected for Familywise Error (FWE).

Diffusion-weighted imaging data were analyzed using the Oxford Centre for Functional Magnetic Resonance Imaging of the Brain (FMRIB) Software Library (FSL; www.fmrib.ox.ac.uk/fsl). DTI data were edited for eddy current-induced image distortions using affine, multi-scale, two-dimensional registration. Fiber tracking was performed using a probabilistic tractography method based on a multifiber model. In the present study we used tractography

routines implemented in the FMRIB Diffusion Toolbox (FDT) (5,000 streamline samples from voxel seeds, 0.5 mm step length, 0.2 curvature threshold)^{17–19}. The seed areas were determined from the fMRI results of each subject (the activated cluster including the precentral knob) (Fig. 1). Of the 5,000 samples generated from each seed voxel, the results for each contact were visualized at a minimum threshold of 25 streamlines¹⁷ and a maximum of 5,000 streamlines per voxel for analysis. Finally, we defined several regions of interest (ROIs) in the brain. The neural network developed from the fMRI results was investigated to determine its incidence in the ROIs. Percentage of incidence (N%) was calculated using the following equation:

$$N\% = \frac{\text{The term of total incidence in each ROI}}{\text{Total case (38 hemispheres)}} \times 100$$

Several ROIs were defined using the diffusion-weighted EPI volume with the highest signal-to-noise ratio (no diffusion weighting, $b=0$) in each subject's brain. We subdivided the frontal cortex into the primary motor cortex (M1); Bordmann area (BA) 4, the premotor cortex (PMC); posterior part of BA 6²⁰, the supplementary motor area (SMA); anterior and medial part of BA 6, the pars opercularis; BA 44; and the pars triangularis, BA 45²¹. The temporal cortex was classified into three gyri, according to morphology: the superior temporal gyrus, middle temporal gyrus, and inferior temporal gyrus. The parietal cortex was subdivided into the primary somatosensory cortex (S1); BA 3, 1, and 2; superior parietal lobule (SPL); intraparietal sulcus (IPS); the supramarginal gyrus^{22,23}; and the angular gyrus¹⁰. The thalamus was subdivided into the lateral group (including the ventral posterior lateral nucleus (VPL)), medial group, posterior group, and anterior group²⁴. The putamen was defined as having anterior, middle, and posterior areas²⁵. The cerebellum was divided into contralateral and ipsilateral hemispheres.

The terms of total incidence in each ROI were used to perform a chi-square test for the determination of variances between the right and left hemispheres. Statistical significance was chosen as $P < 0.05$.

RESULTS

The contralateral SM1s are activated during movements of either hand in all subjects. Therefore, we used the activation of SM1 as a seed for analysis of the neural network for hand movement. We defined a successful result as a connection with SM1 that was observed in more than half of the total number of hemispheres (all subjects) in each ROI. In the frontal lobe activations of the contralateral SM1 were connected with the contralateral PMC (100%), contralateral SMA (89.47%), contralateral pars opercularis (81.58%), contralateral pars triangularis (68.42%), and ipsilateral M1 (60.53%). These were also connected with the contralateral SPL (100%), contralateral IPS (100%), contralateral supramarginal gyrus (97.37%), and contralateral angular gyrus (65.79%) in the parietal lobe,

Table 1. Percentage of tracks that reached the region of interest

Regions of interest	Total (%)
Contralateral premotor cortex (PMC) (BA 6)	100
Contralateral superior parietal lobule (SPL) (BA 5,7)	100
Contralateral intraparietal sulcus (IPS)	100
Contralateral supramarginal gyrus (BA40)	97.37
Contralateral supplementary motor area (SMA) (BA 6)	89.47
Contralateral middle temporal gyrus (BA 21)	89.47
Contralateral VPL nucleus of thalamus	86.84
Contralateral posterior one-third of putamen	81.58
Contralateral pars opercularis (BA 44)	81.58
Contralateral superior temporal gyrus (BA 41,42,22)	71.05
Contralateral pars triangularis (BA 45)	68.42
Contralateral angular gyrus (BA 39)	65.79
Ipsilateral M1 (BA 4)	60.53
Contralateral cerebellum hemisphere	60.53

Total (%) indicates percentage of the tracks that reached anatomical areas in both hemispheres.

and the contralateral superior temporal gyrus (71.05%) and contralateral middle temporal gyrus (89.47%) in the temporal lobe. The contralateral VPL nucleus of the thalamus (86.84%), contralateral posterior one-third of the putamen (81.58%), and contralateral hemisphere of the cerebellum (60.53%) were connected with the ipsilateral SM1 activation region (Table 1) (Fig 1). There was no significant difference in the total ROI incidence between the right and left hemispheres ($p>0.05$).

DISCUSSION

In the current study we elucidated the neural network for hand movement using a combined fMRI/DTT method. Our results show that SM1 connected to the PMC (100%), SPL (100%), IPS (100%), supramarginal gyrus (97.37%), SMA (89.47%), VPL nucleus of the thalamus (86.84%), posterior one-third of the putamen (81.58%), pars opercularis (81.58%), pars triangularis (68.42%), angular gyrus (65.79%), and cerebellum (60.53%) in the same hemisphere and the contralateral primary motor cortex (60.53%) in the opposite hemisphere. The classical conception of motor control holds that the corticospinal tract (CST) is the most important motor pathway for voluntary movements in humans. The majority of CSTs originate from pyramidal neurons, which are situated in SM1^{26,27}. In this study the contralateral SM1 was activated during hand movements of either hand in all subjects. Previous neurophysiology studies have reported that neurons of the PMC, SMA, and cerebellum are involved in the control of movement velocity, amplitude, and directional tuning with SM1^{28–31}. It is widely accepted that motor areas receive afferents from a specific set of parietal areas^{32,33}. Several researchers have demonstrated that the SPL- and IPS-integrated somatosensory and visual information participate in goal-direction actions^{32–34}. Other neuroimaging studies have provided evidence that these areas are involved in the praxis along with the SMA, PMC, and cerebellum^{35–39}. In addition to these areas, linearly increasing activations corresponding to movement amplitude and velocity were observed in the putamen and thalamus^{3,35,39}. The supramarginal gyrus and

angular gyrus areas are concerned with motor planning and cognitive judgments, such as mechanical problem solving^{22,40,41}). For example, increase of neural activation in the supramarginal gyrus and angular gyrus is based on preparation of pointing movements. This cortical recruitment is related to the more complex coordinate transformations needed to plan a pointing movement²²). Rao et al. verified that the pars opercularis and pars triangularis connections were activated more substantially during the performance of the complex task than during the simple task⁴²). According to other studies the pars opercularis and pars triangularis were activated by imitation and observation of finger movements^{43,44}). Therefore, these areas seem to constitute a high-level sensorimotor interface integrating sensory stimuli and cognitive tasks with the related motor representations of hand movements⁴⁵). We also found connections to the ipsilateral M1 in the opposite hemisphere. Many previous studies have demonstrated the contribution of the ipsilateral M1 to motor control in normal portions of the injured brain^{36,46,47}). We observed unexpected connections to the middle and superior temporal gyri. These areas are involved in the recognition of an object²³); therefore, there may be potential partial volume contamination in the centrum semiovale⁴⁸).

There have been many neuroimaging and neurophysiologic studies of neural control of hand movement, and our results generally coincide with those of previous studies^{3,28–31,34–39}). To the best of our knowledge, only one DTT study has addressed this topic. Guye et al. investigated the connectivity of M1 using combined fMRI and DTI¹²). They implemented fast-marching tractography to define connectivity maps within the whole brain, and the connectivity maps were normalized. The results showed strong connections from M1 to the CST, premotor areas, parietal cortices, thalamus, and cerebellum. In contrast, we used probabilistic tractography to analyze DTI data. Probabilistic tractography, which is a multi-tensor model, estimates multiple dominant diffusion orientations within an imaging voxel^{10,18,48}). Behrens et al. reported that probabilistic tractography was more sensitive than traditional tensor methods or single-fiber approaches¹⁷). We found that SM1 is connected with brain areas, such as the supramarginal gyrus, angular gyrus, pars opercularis, pars triangularis, and contralateral M1, in addition to the areas previously identified in the study of Guye et al.¹²).

In conclusion, we visualized the neural network related to hand movements and found this neural network consisted of many brain areas in addition to those previously identified for motor planning and execution. We believe the results and methodology of this study will be helpful in the investigation of motor control in the human brain. Limitations of the present study included the false positives generated by the multi-fiber approach due to the inability of DTI to fully reflect the underlying fiber architecture because of regions of fiber complexity and crossing^{17,25}). A clear understanding of the relationship between fiber bundle structure and DTI will allow us to elucidate a more accurate neural network for movements.

ACKNOWLEDGEMENT

This work was supported by National Research Foundation of Korea Grant funded by the Korean Government (KRF-2008-314-E00173).

REFERENCE

- 1) Schieber MH, Santello M: Hand function: peripheral and central constraints on performance. *J Appl Physiol* 2004, 96: 2293–2300.
- 2) Eickhoff SB, Dafotakis M, Grefkes C, et al.: Central adaptation following heterotopic hand replantation probed by fMRI and effective connectivity analysis. *Exp Neurol* 2008, 212: 132–144.
- 3) Kim JA, Eliassen JC, Sanes JN: Movement quantity and frequency coding in human motor areas. *J Neurophysiol* 2005, 94: 2504–2511.
- 4) Meindl T, Born C, Britsch S, et al.: Functional BOLD MRI: comparison of different field strengths in a motor task. *Eur Radiol* 2008, 18: 1102–1113.
- 5) Taniguchi M, Kato A, Fujita N, et al.: Movement-related desynchronization of the cerebral cortex studied with spatially filtered magnetoencephalography. *Neuroimage* 2000, 12: 298–306.
- 6) Urbano A, Babiloni C, Onorati P, et al.: Dynamic functional coupling of high resolution EEG potentials related to unilateral internally triggered one-digit movements. *Electroencephalogr Clin Neurophysiol* 1998, 106: 477–487.
- 7) Wang C, Wai Y, Kuo B, et al.: Cortical control of gait in healthy humans: an fMRI study. *J Neural Transm* 2008, 115: 1149–1158.
- 8) Bassar PJ, Mattiello J, LeBihan D: Estimation of the effective self-diffusion tensor from the NMR spin echo. *J Magn Reson B* 1994, 103: 247–254.
- 9) Bassar PJ, Jones DK: Diffusion-tensor MRI: theory, experimental design and data analysis - a technical review. *NMR Biomed* 2002, 15: 456–467.
- 10) Rushworth MF, Behrens TE, Johansen-Berg H: Connection patterns distinguish 3 regions of human parietal cortex. *Cereb Cortex* 2006, 16: 1418–1430.
- 11) Kim SG, Ashe J, Hendrich K, et al.: Functional magnetic resonance imaging of motor cortex: hemispheric asymmetry and handedness. *Science* 1993, 261: 615–617.
- 12) Guye M, Parker GJ, Symms M, et al.: Combined functional MRI and tractography to demonstrate the connectivity of the human primary motor cortex in vivo. *Neuroimage* 2003, 19: 1349–1360.
- 13) Jang SH, Ahn YH, Kim SH, et al.: Corticospinal tract restoration: combined study of diffusion tensor tractography, functional MRI, and transcranial magnetic stimulation. *J Comput Assist Tomogr* 2007, 31: 901–904.
- 14) Kim YH, Kim DS, Hong JH, et al.: Corticospinal tract location in internal capsule of human brain: diffusion tensor tractography and functional MRI study. *Neuroreport* 2008, 19: 817–820.
- 15) Oldfield RC: The assessment and analysis of handedness: the Edinburgh inventory. *Neuropsychologia* 1971, 9: 97–113.
- 16) Cherubini A, Luccichenti G, Peran P, et al.: Multimodal fMRI tractography in normal subjects and in clinically recovered traumatic brain injury patients. *Neuroimage* 2007, 34: 1331–1341.
- 17) Behrens TE, Berg HJ, Jbabdi S, et al.: Probabilistic diffusion tractography with multiple fibre orientations: What can we gain? *Neuroimage* 2007, 34: 144–155.
- 18) Behrens TE, Woolrich MW, Jenkinson M, et al.: Characterization and propagation of uncertainty in diffusion-weighted MR imaging. *Magn Reson Med* 2003, 50: 1077–1088.
- 19) Owen SL, Heath J, Kringelbach M, et al.: Pre-operative DTI and probabilistic tractography in four patients with deep brain stimulation for chronic pain. *J Clin Neurosci* 2008, 15: 801–805.
- 20) Picard N, Strick PL: Imaging the premotor areas. *Curr Opin Neurobiol* 2001, 11: 663–672.
- 21) Dronkers NF, Plaisant O, Iba-Zizen MT, et al.: Paul Broca's historic cases: high resolution MR imaging of the brains of Leborgne and Lelong. *Brain* 2007, 130: 1432–1441.
- 22) Astafiev SV, Shulman GL, Stanley CM, et al.: Functional organization of human intraparietal and frontal cortex for attending, looking, and pointing. *J Neurosci* 2003, 23: 4689–4699.
- 23) Ramayya AG, Glasser MF, Rilling JK: A DTI Investigation of Neural Substrates Supporting Tool Use. *Cereb Cortex* 2010, 20: 507–516.
- 24) Morel A, Magnin M, Jeanmonod D: Multiarchitectonic and stereotactic atlas of the human thalamus. *J Comp Neurol* 1997, 387: 588–630.
- 25) Lehericy S, Ducros M, Krainik A, et al.: 3-D diffusion tensor axonal tracking shows distinct SMA and pre-SMA projections to the human striatum. *Cereb Cortex* 2004, 14: 1302–1309.

- 26) Afifi AK, Bergman RA: Functional neuroanatomy : text and atlas (2nd ed). New York: Lange Medical Books/McGraw-Hill, 2005, pp 232–235.
- 27) Georgopoulos AP, Kalaska JF, Caminiti R, et al.: On the relations between the direction of two-dimensional arm movements and cell discharge in primate motor cortex. *J Neurosci* 1982, 2: 1527–1537.
- 28) Fu QG, Flament D, Coltz JD, et al.: Temporal encoding of movement kinematics in the discharge of primate primary motor and premotor neurons. *J Neurophysiol* 1995, 73: 836–854.
- 29) Fu QG, Mason CR, Flament D, et al.: Movement kinematics encoded in complex spike discharge of primate cerebellar Purkinje cells. *Neuroreport* 1997, 8: 523–529.
- 30) Fu QG, Suarez JJ, Ebner TJ: Neuronal specification of direction and distance during reaching movements in the superior precentral premotor area and primary motor cortex of monkeys. *J Neurophysiol* 1993, 70: 2097–2116.
- 31) He SQ, Dum RP, Strick PL: Topographic organization of corticospinal projections from the frontal lobe: motor areas on the lateral surface of the hemisphere. *J Neurosci* 1993, 13: 952–980.
- 32) Rizzolatti G, Luppino G, Matelli M: The organization of the cortical motor system: new concepts. *Electroencephalogr. Clin Neurophysiol* 1998, 106: 283–296.
- 33) Rizzolatti G, Luppino G: The cortical motor system. *Neuron* 2001, 31: 889–901.
- 34) Caminiti R, Ferraina S, Johnson PB: The sources of visual information to the primate frontal lobe: a novel role for the superior parietal lobule. *Cereb Cortex* 1996, 6: 319–328.
- 35) Agnew JA, Zeffiro TA, Eden GF: Left hemisphere specialization for the control of voluntary movement rate. *Neuroimage* 2004, 22: 289–303.
- 36) Chang MC, Ahn SH, Cho YW, et al.: The comparison of cortical activation patterns by active exercise, proprioceptive input, and touch stimulation in the human brain: A functional MRI study. *NeuroRehabilitation* 2009, 25: 87–92.
- 37) Choi SH, Na DL, Kang E, et al.: Functional magnetic resonance imaging during pantomiming tool-use gestures. *Exp Brain Res* 2001, 139: 311–317.
- 38) Turner RS, Grafton ST, Votaw JR, et al.: Motor subcircuits mediating the control of movement velocity: a PET study. *J Neurophysiol* 1998, 80: 2162–2176.
- 39) Turner RS, Desmurget M, Grethe J, et al.: Motor subcircuits mediating the control of movement extent and speed. *J Neurophysiol* 2003, 90: 3958–3966.
- 40) Buxbaum LJ: Ideomotor apraxia: a call to action. *Neurocase* 2001, 7: 445–458.
- 41) Johnson-Frey SH, Newman-Norlund R, Grafton ST: A distributed left hemisphere network active during planning of everyday tool use skills. *Cereb Cortex* 2005, 15: 681–695.
- 42) Rao H, Di X, Chan RC, et al.: A regulation role of the prefrontal cortex in the fist-edge-palm task: evidence from functional connectivity analysis. *Neuroimage* 2008, 41: 1345–1351.
- 43) Binkofski F, Amunts K, Stephan KM, et al.: Broca's region subserves imagery of motion: a combined cytoarchitectonic and fMRI study. *Hum Brain Mapp* 2000, 11: 273–285.
- 44) Iacoboni M, Woods RP, Brass M, et al.: Cortical mechanisms of human imitation. *Science* 1999, 286: 2526–2528.
- 45) Binkofski F, Buccino G: Motor functions of the Broca's region. *Brain Lang* 2004, 89: 362–369.
- 46) Grefkes C, Nowak DA, Eickhoff SB, et al.: Cortical connectivity after subcortical stroke assessed with functional magnetic resonance imaging. *Ann Neurol* 2008, 63: 236–246.
- 47) Li A, Yetkin FZ, Cox R, et al.: Ipsilateral hemisphere activation during motor and sensory tasks. *AJNR Am J Neuroradiol* 1996, 17: 651–655.
- 48) Parker GJ, Alexander D: Probabilistic anatomical connectivity derived from the microscopic persistent angular structure of cerebral tissue. *Philos Trans R Soc Lond B Biol Sci* 2005, 360: 893–902.